

# Comparison of autotrophic and mixotrophic cultivation of green microalgal for biodiesel production

Yu-Ru Li<sup>1</sup>, Wen-Tien Tsai<sup>1</sup>, Yi-Chyun Hsu<sup>2</sup>, Meng-Zhi Xie<sup>3</sup>, Jen-Jeng Chen<sup>4,\*</sup>

<sup>1</sup>Graduate Institute of Bioresources, National Pingtung University of Science and Technology, Pingtung 91201, Taiwan

<sup>2</sup>Department of Environmental Engineering, Kun Shan University, Tainan 71003, Taiwan

<sup>3</sup>Department of Pharmacy and Graduate Institute of Pharmaceutical Technology, Tajen University, Pingtung 90741, Taiwan

<sup>4</sup>Graduate Institute of Environmental Management, Tajen University, Pingtung 90741, Taiwan

## ABSTRACT

The effects of autotrophic and mixotrophic growth on cell growth and lipid productivity of green microalga *Chodatella* sp. were investigated. Carbon dioxide and piggery wastewater served as the carbon and nutrient sources, respectively, for autotrophic and mixotrophic growth. Appropriate doses of each source were found to be beneficial to biomass production. The cultures produced similar fatty acid compositions, which are suitable for biodiesel production. The specific growth rate, biomass production, and lipid productivity obtained with mixotrophic growth were 1.74, 14, and 5.6 times higher than those obtained with autotrophic growth, respectively. The mixotrophic cultivation simultaneously assimilated 99.7% ammonia nitrogen and 75.9% total phosphorus from piggery wastewater, which reduced the required nutrient for the culture of microalgae, thereby reducing the cost of biodiesel production.

**Keywords:** autotrophic, biodiesel production, mixotrophic

## INTRODUCTION

Despite many microalgae being exceedingly rich in lipid content as biodiesel feedstock, limitations such as efficacy and cost-competitiveness must be addressed [1, 2]. One of the methods used for reducing the cost of algal biomass is to integrate wastewater treatment with algae biomass production [3]. For example, Lau et al. [4] reported that over 90% of N content and 80% of P content were removed by *Chlorella vulgaris* from primary treated sewage. A similar microalgal strain, *Chlorella pyrenoidosa*, in soybean processing wastewater was demonstrated not only to remove 78% of soluble organic carbon, 89% of total nitrogen (TN), and 70% of total phosphate (TP), but also attained an average biomass productivity of 0.64 g/L·d with an average lipid content of 37% [5]. Sydney et al. [6] found that *Botryococcus braunii* is able to remove N and P nutrients (79.63%) from treated domestic wastewater, and accumulate oil with a dry biomass of up to 36%. The lipid profiles of this extracted oil are similar to those of oilseed feedstocks.

The accumulation of lipids and the cell growth rate in wastewater depend on diverse factors, such as cultivation conditions, algal species, and growth environment [1,3]. An excessive nutrient concentration can inhibit algal

growth [7,8]. The organic carbon/nitrogen (C/N) ratio and the carbon/phosphorous (C/P) ratio in wastewater may be low compared to the typical ratios in rapidly growing algal biomass (C/N 6:1; C/P 48:1) [8]. Li et al. [9] found that at a 5-8:1 N/P ratio, both nutrients can be highly removed by *Scenedesmus* sp. in an artificial medium. Li et al. [9] also found that N or P limitation can lead to the accumulation of lipids (30% vs. 53%). However, there is a trade off between maximal algal lipid content and optimal algal growth in wastewater, the former requiring low nutrient content and the latter requiring high organic nutrient concentration [10]. A major limitation is that despite inducing very high lipid content, the cell growth rate is often very low, which limits lipid production [11,12].

Swine breeding is an important industry in Taiwan, with piggery wastewater discharged from over 6 millions pigs. Algal growth in wastewater that contributes to renewable energy production and wastewater treatment is a promising strategy. However, there is little information available for cultivating microalgae in a piggery wastewater environment. The strain of *Chodatella* sp. isolated from local fresh water was found to grow in both autotrophic and mixotrophic modes. The present study compares the cell growth rate and biodiesel production under autotrophic and mixotrophic growth.

## MATERIALS AND METHODS

### 2.1 Microalgal culturing

One of the dominant green microalgal species, *Chodatella* sp., was isolated from local source water and cultured in a medium according to the method presented by Norris et al. [13]. Autotrophic cultures of *Chodatella* sp. were grown in batch mode in a 5-L modified serum bottle containing 4 L of sterilized algal medium. The bottles were placed in an incubator maintained at 25 °C and continuously provided with 5 klux of illumination. CO<sub>2</sub> was supplied to the cultures by syringe injection every day. For mixotrophic growth, cells were incubated with piggery wastewater at 25 °C with continuous illumination of 5 klux and aeration with air at a 0.8 L/min·L flow rate. The piggery wastewater effluent was collected from a local pig farm at Pingtung University of Science and Technology, Taiwan. The piggery wastewater was filtered through a 0.45-μm membrane and sterilized before experiments. Cultures were harvested in the log growth phase for experiments. For autotrophic cultures,

Corresponding Author: [jerry@tajen.edu.tw](mailto:jerry@tajen.edu.tw)

the CO<sub>2</sub> concentrations of the medium were 2, 4, and 8% (vol.), and the piggery wastewater content were 20, 50, and 100% for mixotrophic cultures, respectively.

## 2.2. Biomass concentration and growth rate

The optical density of microalgal cells was determined by measuring absorbance at 684 nm (OD<sub>684</sub>) using an ultraviolet/visible spectrophotometer (Model U-2001, Hitachi, Japan). The dry weight of the microalgal biomass was determined gravimetrically. A known volume of microalgal culture was collected and dried at 90 °C for 3 hours. The growth rate ( $\mu$ ) was calculated according to the equation  $\mu = (\ln A_1 - \ln A_0)/(T_1 - T_0)$ , where A1 and A0 are the dry weights of the microalgal biomass at times T1 and T0, respectively. The relationship between microalgal biomass and OD<sub>684</sub> is: biomass = 14.447 + 259.678 OD<sub>684</sub> ( $R^2 = 0.972$ ,  $P < 0.0001$ ).

## 2.3. Lipid content

A stock culture of *Chodatella* sp. cells was collected by centrifugation at 5000 × g for 5 min. The precipitated algal cells were washed and resuspended in deionized water in triplicate. Cells were collected by centrifugation again and then dried by a freeze dryer. The microalgal total lipids were extracted with n-hexane/methanol (2/1, v/v) in a Soxhlet extractor and quantified gravimetrically. The biomass concentration (mg/L) is expressed as the dry weight of the microalgal biomass.

## 2.4. Fatty acid methyl esters

Freeze-dried biomass (0.1 g) was placed in 50-mL Teflon-capped Pyrex tubes and mixed with a premixed homogeneous solution of NaOH catalyst (2.5 wt.%) and methanol. 8.0 mL of alkali catalyst was added to the tubes. The reaction mixture was heated at 60 °C for 30 min, with the samples well-mixed during heating. After the reaction was completed, the tubes were allowed to cool to room temperature. Then, 8 mL of an acid catalyst (HCl:methanol, 5.8 vol.%) and 10 mL of a boron trifluoride/methanol solution were added to the tubes in a water bath at 100 °C for 15 min. After the reaction was completed, the tubes were allowed to cool to room temperature. Purification of the solution was achieved by using 2 mL of saturated sodium chloride solution. 4 mL of an extracting solvent (hexane) was then added to each tube. The upper solvent layer containing fatty acid methyl esters (FAMES) was collected and analyzed by gas chromatography.

FAME analysis was performed using a gas chromatograph with an autosampler (Agilent 7820A, USA), flame ionization detector, and a DB-23 Agilent column (length: 60 m, ID: 0.25 mm, film: 0.25  $\mu$ m). Hydrogen was used as the carrier gas at a constant flow rate of 35 mL/min. The injector was held at 270 °C while the detector was kept at 280 °C. The oven was at 130 °C initially, ramped to 170 °C at 6.5 °C/min, then ramped again to 215 °C at 2.75 °C/min, and held at 215 °C for 12 min. The temperature was further increased to 230 °C at a heating rate of 40 °C/min and held for 3 min, giving a total heating time of 38.88 min. FAMES were identified

by comparing their retention times with those of standard fatty acids (Supelco, USA) and quantified by comparison with the prepared calibration curves. Pentadecanoic acid (C15:0) was used as an internal standard.

## 2.5. Nitrogen and phosphorus analysis

The samples were first filtered through a 0.45- $\mu$ m membrane and then the filtrate was properly diluted and analyzed for ammonia nitrogen (NH<sub>3</sub>-N), organic nitrogen, and orthophosphate (PO<sub>4</sub><sup>3-</sup>-P). NH<sub>3</sub>-N and PO<sub>4</sub><sup>3-</sup>-P concentrations were determined according to Standard Methods [14]. Total Kjeldahl nitrogen (TKN), the sum of NH<sub>3</sub>-N and organic nitrogen, was measured in the experiments.

## RESULTS AND DISCUSSION

### 3.1 Effect on cell growth

The cell growth indicated by optical density for various CO<sub>2</sub> and piggery wastewater content levels is shown in Fig. 1. It can be seen that *Chodatella* sp. survived in all of the cultures and no obvious lag phases were observed except for the culture under 100% piggery wastewater content. A comparison of the cultures shows that cells grew faster under piggery wastewater cultivation than under CO<sub>2</sub> cultivation. For example, the optical density of the cells was 1.87 cm<sup>-1</sup> at 100% piggery wastewater content in the 7<sup>th</sup> day. Under CO<sub>2</sub> cultivation, the optical density of the cells reached 0.693 cm<sup>-1</sup> at 4% content in the same cultivation period. The average specific growth rates under cultures with piggery wastewater content levels of 20, 50, and 100% were 0.24, 0.281, and 0.303 d<sup>-1</sup>, respectively. These specific growth rates are higher than those under CO<sub>2</sub> cultivation (0.14, 0.173, and 0.114 d<sup>-1</sup> for CO<sub>2</sub> concentrations of 2, 4, and 8%, respectively).

Several factors influence algal cell growth, including the availability of nutrients, light (quality and quantity), pH, temperature, and the initial inoculation density. Figure 1a shows the growth curve under CO<sub>2</sub> cultivation. The optimal CO<sub>2</sub> content for cell growth is 4%. Insufficient CO<sub>2</sub> concentration (2%) in the culture would not sustain optimal growth. However, higher CO<sub>2</sub> concentration did not further improve cell growth. Similar results were reported for *Scenedesmus obliquus* by Kaewkannetra et al. [15]. This is probably due to excess CO<sub>2</sub> being converted to H<sub>2</sub>CO<sub>3</sub>, resulting in a reduction of pH in the culture, thereby affecting cell growth. Greque de Moraes and Costa [16] reported that a *Spirulina* sp. culture with 6% CO<sub>2</sub> had the maximum specific growth rate. Higher CO<sub>2</sub> concentration led to a decrease of biomass yield.

Under piggery wastewater cultivation, the optimal content for cell growth is 50% (Fig. 1b). With 100% piggery wastewater content, the culture had a lag phase during the first 2 days. It is speculated that this may be due to the relatively high ratio of nutrient concentration to initial inoculation density. Cell growth dramatically increased after 4 days under 100% piggery wastewater content. This mixotrophic culture produced a 1.74 times

higher specific growth rate compared with that for autotrophic growth with 4% CO<sub>2</sub> content.




Fig. 1. Cell growth in various content of (a) CO<sub>2</sub> and (b) piggery wastewater cultivation

### 3.2 Effect on biomass and lipid productivity

Fig. 2 shows the effects of CO<sub>2</sub> and piggery wastewater cultivation on biomass and lipid productivity. Although both CO<sub>2</sub> and piggery wastewater cultivations could sustain microalgal growth, the biomass and lipid productivity obtained with piggery wastewater were much higher than those obtained with CO<sub>2</sub>. Even with the lowest productivity for piggery wastewater cultivation (20% content), biomass and lipid productivity were higher than those obtained with the optimal CO<sub>2</sub> content (4%). The biomass productivity (Fig. 2c) reached up to 115 mg/L·d under 50% piggery wastewater content, whereas that with cultivation with 4% CO<sub>2</sub> content was only 28.6 mg/L·d (Fig. 2a). Biomass production and the associated cell growth, as demonstrated by biomass productivity, indicate that mixotrophic growth with piggery wastewater is more beneficial than autotrophic growth. This result is similar to the findings for *Chlorella vulgaris* [17] and *Chlorella protothecoides* [18], where mixotrophic cultures with glucose or acetate had higher biomass productivities than those of autotrophic cultures.

Lipid productivity is the mass of lipids produced per unit volume of the culture per unit time; it depends on the algal growth rate and the lipid content of the biomass. Fig. 2b shows the effect of CO<sub>2</sub> content on lipid productivity. The highest lipid productivity was achieved with 4% CO<sub>2</sub>

content, implying that there exists an optimal dosage for CO<sub>2</sub> cultivation. This finding is consistent with the biomass productivity data. However, 8% CO<sub>2</sub> content resulted in a lower lipid content in cells (data not shown); therefore, a decrease in lipid productivity is more significant than that of biomass productivity. The results show that high CO<sub>2</sub> content not only decreased cell growth, but also significantly hindered lipid accumulation.

Under piggery wastewater cultivation, the biomass productivity increased when the content was higher than 50%. The culture with 100% wastewater produced a lower lipid content than that produced by the culture with 50% wastewater (Fig. 2d), and thus had lower lipid productivity. The 50% piggery wastewater content culture reached 51.4 mg/L·d lipid productivity, approximately 14 times higher than that obtained with the culture with 4% CO<sub>2</sub> content. The better performance of mixotrophic growth with piggery wastewater compared to that of autotrophic growth is conjectured to be due to the homogeneous distribution of nutrients and favorable carbon utilization in mixotrophic culture. This lipid productivity is much higher than that reported by Wang [19] for *Chlorella pyrenoidosa* cultivated in piggery wastewater (maximum lipid productivity of 6.3 mg/L·d).




Fig. 2. Biomass and lipid productivity in various content of (a, b) CO<sub>2</sub> and (c, d) piggery wastewater cultivation

### 3.3 Effect on fatty acid content and composition

Fig. 3 compares the FAME concentration and FAME composition of cultures with the optimal levels of CO<sub>2</sub> and piggery wastewater. Similar to lipid productivity, the amount of total FAMES produced from piggery wastewater cultivation was much higher than that obtained with CO<sub>2</sub> cultivation. Under the testing conditions, cultivation with 4% CO<sub>2</sub> and 50% piggery wastewater produced similar fatty acid compositions, comprising mainly palmitic acid (C16:0), oleic acid (C18:1), linoleic acid (C18:2), and linolenic acid (C18:3). Oleic acid (C18:1) was the most abundant fatty acid in algal cells under piggery wastewater cultivation. Palmitic acid (C16:0) was the second most abundant fatty acid. Palmitic and oleic and fatty acids were also the most abundant fatty acids for CO<sub>2</sub> cultivation.

These four major fatty acids produced from autotrophic CO<sub>2</sub> growth and mixotrophic piggery wastewater growth are quite similar to those of soybean oil and rapeseed oil and are thus suitable for biodiesel production. However, the total amount of FAMES obtained with mixotrophic growth was 5.6 times higher than that obtained with autotrophic growth. This indicates that lipids produced by microalgae under mixotrophic growth have great potential as economical feedstock for biodiesel production, due to the fact that piggery wastewater can serve the essential nutrients and carbon supply for algae growth. Sturm et al. [20] confirmed that algal biofuel production is energetically favorable when utilizing wastewater as a nutrient source. Mixotrophic growth produces algal biomass while removing nutrients from wastewater, making it a feasible and promising strategy [21].





Fig. 3. Fatty acid concentration and compositions of (a) CO<sub>2</sub> and (b) piggery wastewater cultivation

### 3.4 Biomass production with nutrient removal

Fig. 4 shows the changes in TN, NH<sub>3</sub>-N, TKN, PO<sub>4</sub><sup>3-</sup>-P, biomass concentration, and cell lipid content with time for cultivation with 50% piggery wastewater. Ammonium decreased from 44.83 mg/L to 0.16 mg/L and TKN dropped from 69.04 mg/L to 40.99 mg/L after 12 days of cultivation (Fig. 4a). The total phosphorus drastically decreased from 2.79 mg/L to 0.67 mg/L in the same period (Fig. 4b). The removal efficiencies of TKN, NH<sub>3</sub>-N, and TP are 48.6%, 99.7%, and 75.9%, respectively. As mentioned above, TKN is the sum of NH<sub>3</sub>-N and organic nitrogen. Organic nitrogen increased after cell growth (data not shown) due to the release of cell endogenous decay. Therefore, ammonium is main form of nitrogen to provide sufficient nutrient for meeting the cell growth in piggery wastewater [22].

The decrease in nitrogen and phosphorus resulted in an increase in biomass concentration (Fig. 4c). The biomass concentration increased from 136 mg/L to 1,406 mg/L after 12 days. The cell lipid content also increased with cultivation time, up to 41.2% at the 12<sup>th</sup> day. Numerous researchers have reported that algal cell lipids accumulate under nutrient limitation in the culture [23-25]. However, nutrient limitation inhibits cell growth, and thus lowers lipid productivity. In this case, no significant reduction of biomass production was observed even under lower NH<sub>3</sub>-N and TP concentrations after the 4<sup>th</sup> day. Specific nutrient removal rates of 3.72 mg N/L·d and 0.18 mg P/L·d were obtained. These results are consistent with the nitrogen and phosphorous removal rates reported by Ruiz et al. [26] for a culture of freshwater *Scenedesmus obliquus* in a synthetic medium (removal rates of 13.5-4.2 mg N/L·d and 1.49-0.32 mg P/L·d).




Fig. 4. (a) Nitrogen concentration, (b) total phosphorus concentration, (c) biomass concentration, and (d) cell lipid content during batch culture on 50% piggery wastewater

## CONCLUSION

The effects of cultivation conditions on algal biomass and lipid production of green microalgae (*Chodatella* sp.) grown in autotrophic and mixotrophic modes were investigated. Carbon dioxide and piggery wastewater served as the carbon and nutrient sources, respectively. Under both autotrophic and mixotrophic cultivation conditions, an appropriate dose of each source was beneficial to biomass production. Both cultivation method led to similar fatty acid compositions, comprising palmitic (C16:0), oleic (C18:1), linoleic (C18:2), and linolenic (18:3) acids. The specific growth rate, biomass production, and FAME yield obtained with mixotrophic growth were much higher than those obtained with autotrophic growth. *Chodatella* sp. is a potential strain for integrating piggery wastewater treatment with algae biomass production. Its mixotrophic growth produced biodiesel practically and economically.

## ACKNOWLEDGMENT

The authors would like to thank the National Science Council of Taiwan for financially supporting this research under grants NSC 99-2221-E-127-004 and NSC 100-2221-E-127-002.

## REFERENCES

- [1] Chisti Y.; *Biodiesel from microalgae*. Biotechnol Adv, vol. 25, 2007, pp. 294-306.
- [2] Brennan L., Owende P.; *Biofuels from microalgae — A review of technologies for production, processing, and extractions of biofuels and co-products*. Renewable and Sustainable Energy Reviews, vol. 24, 2010, pp. 557-577.
- [3] Pittman JK., Dean AP., Osundeko O.; *The potential of sustainable algal biofuel production using wastewater resources*. Bioresour Technol, vol. 102, 2011, pp. 17-25.
- [4] Lau PS., Tam NFY., Wong YS.; *Effect of algal density on nutrient removal from primary settled wastewater*. Environ Pollut, vol. 89, 1995, pp. 59-66.
- [5] Su HY., Zhang YL., Zhang C., Zhou XF., Li JP.; *Cultivation of Chlorella pyrenoidosa in soybean processing wastewater*. Bioresour Technol, vol. 102, 2011, pp. 9884-9890.
- [6] Sydney EB., da Silva TE., Tokarski A., Novak AC., de Carvalho JC., Woiciechowski AL., Larroche C., Soccol CR.; *Screening of microalgae with potential for biodiesel production and nutrient removal from treated domestic sewage*. Appl Energy, vol. 88, 2011, pp. 3291-3294.
- [7] Kong QX., Li L., Martinez B., Chen P., Ruan R.; *Culture of Microalgae Chlamydomonas reinhardtii in Wastewater for Biomass Feedstock Production*. Appl Biochem Biotechnol, vol. 160, 2010, pp. 9-18.
- [8] Metcalf and Eddy.; *Wastewater engineering: Treatment and reuse*. 4th ed. New York: McGraw-Hill; 2003.
- [9] Li X., Hu HY., Gan K., Sun YX.; *Effects of different nitrogen and phosphorus concentrations on the growth, nutrient uptake, and lipid accumulation of a freshwater microalga Scenedesmus sp.* Bioresour Technol, vol. 101, 2010, pp. 5494-5500.
- [10] Abielovich A.; *Water Pollution and Bioremediation by Microalgae*. In: Richmond A, editor. Handbook of microalgal culture, Oxford: Blackwell publishing Inc; 2004, pp. 430-438.



- [11] Scragg AH., Illman AM., Carden A., Shales SW.; *Growth of microalgae with increased caloric values in a tubular bioreactor*. Biomass Bioenerg, vol. 23, 2002, pp. 67-73.
- [12] Lynn SG., Kilham SS., Kreeger DA., Interlandi SJ.; *Effect of nutrient availability on the biochemical and elemental stoichiometry in freshwater diatom Stephanodiscus minutulus (Bacillariophyceae)*. J Phycol, vol. 36, 2000, pp. 510-522.
- [13] Norris L., Norris RE., Calvin M.; *A survey of the rates and products of short-term photosynthesis in plants of nine Phyla*. J of Exp Bot, vol. 6, 1995, pp. 64-74.
- [14] APHA., AWWA., WEF.; *Standard Methods for the Examination of Water and Wastewater*. 19th ed. Washington, DC: American Public Health Association, American Water Works Association and Water Environment Federation; 1995.
- [15] Kaewkannetra P., Enmak P., and Chiu TY.; *The Effect of CO<sub>2</sub> and Salinity on the Cultivation of Scenedesmus obliquus for Biodiesel Production*. Biotechnology and Bioprocess Engineering, vol. 17, 2012, pp.591-597.
- [16] Greque de Moraes M., Costa JAV.; *Biofixation of carbon dioxide by Spirulina sp. and Scenedesmus obliquus cultivated in a three-stage serial tubular photobioreactor*. Journal of Biotechnology, vol. 129, 2007, pp. 439-445.
- [17] Liang Y., Sarkany N., Cui Y.; *Biomass and lipid productivities of Chlorella vulgaris under autotrophic, heterotrophic and mixotrophic growth conditions*. Biotechnol Lett, vol. 31, 2009, pp. 1043-1049.
- [18] Heredia-Arroyo T., Wei W., Hu B.; *Oil Accumulation via Heterotrophic/Mixotrophic Chlorella protothecoides*. Appl Biochem Biotechnol, vol.162, 2010, pp. 1978-1995.
- [19] Wang H., Xiong H., Hui ZL., Zeng XB.; *Mixotrophic cultivation of Chlorella pyrenoidosa with diluted primary piggy wastewater to produce lipids*. Bioresour Technol, vol. 104, 2012, pp. 215-220.
- [20] Sturm BSM., Lamer SL.; *An energy evaluation of coupling nutrient removal from wastewater with algal biomass production*. Applied Energy, vol. 88, 2011, pp. 3499-3506.
- [21] Zhao GL., Yu JY., Jiang FR., Zhang X., Tan TW.; *The effect of different trophic modes on lipid accumulation of Scenedesmus quadricauda*. Bioresource Technology, vol. 114, 2012, pp. 466-471.
- [22] Strett PJ.; Nitrogen assimilation.; In: Lewin RA, editor. *Physiology and biochemistry of algae*. New York: Academic press; 1962, pp. 171-183.
- [23] Yeh KL., Chang JS.; *Effects of cultivation conditions and media composition on cell growth and lipid productivity of indigenous microalga Chlorella vulgaris ESP-31*. Bioresource Technology, vol. 105, 2012, pp. 120-127.
- [24] McGinnis KM., Dempster TA. & Sommerfeld MR.; *Characterization of the growth and lipid content of the diatom Chaetoceros muelleri*. Journal of Applied Phycology 9, 1997, pp. 19-24.
- [25] Lynn SG., Kilham SS., Kreeger DA., Interlandi SJ.; *Effect of nutrient availability on the biochemical and elemental stoichiometry in freshwater diatom Stephanodiscus minutulus (Bacillariophyceae)*. J. Phycol. Vol. 36, 2000, pp. 510-522.
- [26] Ruiz J., Álvarez-Díaz PD., Arbib Z., Garrido-Pérez C., Barragán J., Perales JA.; *Performance of a flat panel reactor in the continuous culture of microalgae in urban wastewater: Prediction from a batch experiment*. Bioresource Technology, vol. 127, 2013, pp. 456-463.